

**PRODUCTION AND CHARACTERIZATION OF BIOMASS AND  
EXOPOLYSACCHARIDES FROM THE EDIBLE MUSHROOM *LENTINUS*  
*SQUARROSULUS***

**MICHAEL T. DADA<sup>1</sup>\*, FAVOUR V. ALABA<sup>1</sup>, FAITH A. BANKOLE<sup>1</sup>, OMOTADE R.  
OGUNREMI<sup>1</sup>**

<sup>1</sup> *Department of Microbiology and Biotechnology, Abiola Ajimobi Technical University, Ibadan 200255, Nigeria*

\* Corresponding author: [michael.dada@tech-u.edu.ng](mailto:michael.dada@tech-u.edu.ng) (Michael T. Dada)

**Abstract:** This study investigated the production and characterization of fungal biomass and exopolysaccharides (EPS) from the indigenous edible wild-type mushroom *Lentinus squarrosulus* (*L. squarrosulus*) grown in submerged fermentation. This mushroom is widely consumed in rural communities but remains underexplored for industrial development of bioproducts. Two nutrient media, Potato dextrose broth (PDB) and Yeast extract broth (YEB) were used for cultivation over 21 days. The biomass yield, EPS concentration and pH changes were measured at intervals of 3 days. The highest EPS ( $0.46 \pm 0.01$  mg/mL) and biomass ( $6.83 \pm 0.043$  mg/mL) production was achieved on day 9 in PDB compared to lesser production with YEB. The nature of EPS was determined by Fourier-transform infrared (FT-IR) spectroscopy that indicated the presence of characteristic functional groups of polysaccharide  $\beta$ -glycosidic bonds. Functional analysis revealed high water solubility ( $54.55 \pm 0.80\%$ ) and water absorbing capacity ( $91.40 \pm 1.20\%$ ), antioxidant assay showed highly active radical scavenging properties ( $87.41 \pm 1.05\%$  equivalent to  $0.415 \pm 0.004$  mg/mL gallic acid). Protein and carbohydrate profiles indicated that EPS was predominantly carbohydrate ( $62.26 \pm 0.16\%$ ). These results show that EPS from *L. squarrosulus* possess favourable structural and bioactive attributes that strongly support its potential use in food-based applications.

**Keywords:** *Lentinus squarrosulus*, exopolysaccharides, submerged fermentation, antioxidant activity, fungal biomass

## RESEARCH ARTICLE

## 1. Introduction

Mushrooms are macrofungi that belong to the phyla Ascomycota and Basidiomycota, together forming the sub-kingdom *Dikarya* (Hibbett *et al.*, 2007). Their rich enzymatic systems enable them to efficiently degrade complex lignocellulosic substrates (Kumla *et al.*, 2020). Although only about 200 out of the estimated 140,000 mushroom species which exist globally are considered edible, and fewer than 25 have been widely cultivated commercially (Stabnikova *et al.*, 2024). This disparity underscores the insufficient utilisation of many indigenous species, which have potential nutritional and industrial values.

Among valuable microbial metabolites, exopolysaccharides (EPS) have attracted considerable interest for industrial applications due to their hydrophilicity, non-toxicity, biocompatibility, and broad functional versatility (Kiran *et al.*, 2024). Diverse biological activities, such as antioxidant, antibacterial, anticancer and immunomodulatory properties of fungal EPS have been shown from numerous studies (Salimi and Farrokh, 2023) and are increasingly considered sustainable alternatives to plant- and seaweed-derived polysaccharides, especially for food, pharmaceutical and cosmetic industries (Abd-Alla *et al.*, 2025).

Structurally, EPS may be categorised as homo-exopolysaccharides (HoEPS), which are made up of a single monosaccharide or hetero-exopolysaccharides (HeEPS), which contain multiple sugar residues (Abd-Alla *et al.*, 2025). It is important to understand their biochemical structure because functional properties such as antioxidant activity, solubility, viscosity and gelling potential are strongly determined by their glycosidic linkages and monosaccharide composition (Wang *et al.*, 2023). As a result, current efforts in research lay emphasis on identifying efficient microbial strains, optimisation of culture conditions and characterisation of the biological and structural attributes of EPS.

Despite the growing interest, most studies on fungal EPS have focused on common mushrooms like *Ganoderma lucidum*, *Lentinula edodes* and *Pleurotus* spp. (Łysakowska *et al.*, 2023). Conversely, many edible tropical mushrooms, including *Lentinus squarrosulus* (*L. squarrosulus*), remain sparsely investigated with respect to biomass composition, EPS yield and functional properties. This knowledge gap is more evident in biodiversity-rich regions such as Nigeria, where there is an abundance of edible mushrooms which are under-utilised due to

## RESEARCH ARTICLE

limited scientific characterisation and dependence on imported bioactive ingredients ([Asonibare et al., 2023](#)).

The present study addresses these knowledge gaps by investigating biomass production and exopolysaccharide yield from *L. squarrosulus* cultivated under submerged fermentation.

This study aims to evaluate the production and characterisation of biomass and EPS from *L. squarrosulus* cultivated under submerged fermentation. The physicochemical and biological properties of the extracted EPS were assessed to determine its potential for food and pharmaceutical applications. It is hypothesised that *L. squarrosulus* produces EPS with favourable physicochemical characteristics and significant antioxidant activity suitable for industrial applications.

## 2. Materials and Methods

### *Culture maintenance*

The macrofungal species *L. squarrosulus* used in this study is a wild-type, indigenous strain originally isolated from a natural habitat and preserved in the culture collection of the Department of Microbiology and Biotechnology, Abiola Ajimobi Technical University, Ibadan, Nigeria. Stock cultures were maintained on Potato dextrose agar (PDA) slants at 4°C.

### *Mycelial transfer and inoculum preparation*

The stock culture of *L. squarrosulus* was subcultured onto freshly prepared PDA plates using a sterile inoculating needle under aseptic conditions. The inoculated plates were incubated at  $28 \pm 2^\circ\text{C}$  for 5 days to obtain a pure culture.

### *Submerged cultivation of *L. squarrosulus* for EPS production*

Actively growing mycelial regions from pure culture were excised using a sterile 1 mm cork borer. These mycelial plugs were then transferred into separate 250 mL Erlenmeyer flasks, each containing 30 mL fresh culture media, which included Yeast extract broth (YEB) and Potato dextrose broth (PDB). Flasks were incubated at  $28 \pm 2^\circ\text{C}$  on a rotary shaker at 120 rpm for 21 days. Sampling was performed at intervals of 3 days by withdrawing one flask for each medium. For each sampling, the pH, EPS production and biomass yield were evaluated using standard procedures. The quantity of the inoculation for each culture medium was 10% (v/v). Submerged fermentation was carried out in the Erlenmeyer flasks for 21 days at 28°C in the two different

## RESEARCH ARTICLE

liquid media under sterile conditions with shaking at 120 rpm ([Ogidi et al., 2020](#)). At intervals of every 3 days, one flask from each medium was sampled to monitor pH, EPS production, and biomass accumulation.

Samples were taken at 3-day intervals for 21 days, pH was determined, and the dry biomass of *L. squarrosulus* was determined by weighing the pellet after centrifugation and oven drying.

***Determination of biomass yield***

Biomass of *L. squarrosulus* was separated from the broth medium by centrifugation at  $4000 \times g$  for 15 min. The resulting mycelial pellet was dried in an oven at  $60^{\circ}\text{C}$  until a constant weight was achieved.

***Determination of pH of culture medium***

The pH of the culture supernatant was measured at every sampling interval using a calibrated digital pH meter.

***Extraction of exopolysaccharides***

Crude exopolysaccharide was precipitated by adding cold 95% ethanol to the culture supernatant in a 1:4 (v/v) ratio and incubating at  $4^{\circ}\text{C}$  overnight to allow complete precipitation of EPS. Exopolysaccharide was collected as a pellet after centrifugation at  $10000 \times g$  for 20 min at  $4^{\circ}\text{C}$ . The resulting pellet was washed with 75% ethanol to remove impurities and centrifuged again at  $10000 \times g$  for 20 min at  $4^{\circ}\text{C}$  ([Ogidi et al., 2020](#)). The EPS pellet was dried in a hot air oven at  $60^{\circ}\text{C}$ , and the weight was recorded as EPS yield in mg/mL.

***Characterization of exopolysaccharides******Fourier transform-infrared (FT-IR) spectroscopy of exopolysaccharides***

The functional groups present in the exopolysaccharide were identified using Fourier transform-infrared (FT-IR) spectroscopy. Two milligrams of EPS powder were ground with 200 mg of dry potassium bromide (KBr) of spectroscopic grade and compressed into a thin pellet. The FT-IR spectrum of the EPS pellet was recorded using a Shimadzu IR Affinity-1S FTIR spectrophotometer (Shimadzu Corporation, Japan) over the spectral range of 4000-400  $\text{cm}^{-1}$  with a resolution of 4  $\text{cm}^{-1}$  at room temperature. The resultant spectrum was analysed to determine the characteristic functional groups and linkages based on their corresponding molecular frequencies. Assigned peaks were cross-referenced with standard FT-IR spectral databases with known absorption patterns, including Sigma-Aldrich IR Spectrum Table version 2019.

## RESEARCH ARTICLE

*Water solubility and water absorption capacity of EPS*

The water solubility index (WSI) and water absorption capacity (WAC) of the crude EPS were determined. Exactly 1.1 mg of each EPS sample was weighed into a pre-weighed centrifuge tube. To this, 1 mL of sterile distilled water was added. The mixture was vortexed and continuously stirred for 30 min at 30°C in a water bath. The suspension was centrifuged at 3000 × g for 10 min to separate the soluble fraction (supernatant) from the insoluble residue (pellet). The supernatant was carefully decanted into a pre-weighed glass Petri dish and dried at 105°C for 4 h in a hot air oven to obtain the weight of dried soluble solids. The remaining wet residue in the centrifuge tube was also weighed (Reddy *et al.*, 2013).

The WSI and WAC were calculated using the following equations:

$$\text{WSI (\%)} = \left( \frac{\text{weight of dry solid in supernatant}}{\text{weight of dry EPS sample}} \right) \times 100$$

$$\text{WAC (\%)} = \left( \frac{\text{weight of wet residue}}{\text{weight of dry EPS sample} - \text{weight of dry solid}} \right) \times 100$$

*Antioxidant activity assay of EPS*

The antioxidant activity of the EPS was performed using the 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging procedure (Gyamfi *et al.*, 1999), with slight modifications. Gallic acid was used as the standard antioxidant. The assay is based on the reduction of DPPH by antioxidants, which results in a change in colour from deep violet to light yellow monitored at 517 nm.

A 0.1 mM DPPH solution was prepared by dissolving it in methanol. Gallic acid standard solutions of different concentrations (1.00, 0.50, 0.25, 0.125 and 0.0313 mg/mL) were obtained by serial dilution in methanol. One milliliter of DPPH solution was mixed with 1 mL of 0.016 mg/mL Exopolysaccharide solution. A solution of methanol and the sample, without DPPH, served as a blank. The control solution was prepared by mixing methanol and the DPPH radical solution. The incubation of all mixtures was carried out for 30 min in the dark, and absorbance was measured at 517 nm using a UV-Visible spectrophotometer.

The percentage DPPH radical scavenging activity (AA%) was determined based on the expression:

## RESEARCH ARTICLE

$$AA\% = \left( \frac{Abs_{Control} - Abs_{Sample}}{Abs_{Control}} \right) \times 100$$

Where, AA% = antioxidant activity (%)

Abs<sub>Sample</sub> = absorbance of the sample

Abs<sub>Control</sub> = absorbance of the control.

The standard curve was generated by plotting the percentage inhibition of gallic acid against concentration. The antioxidant capacity of the EPS was expressed as gallic acid antioxidant capacity (GAEAC) using the equation of the trendline obtained from the graph.

#### *Determination of protein content of EPS*

The protein content of the EPS was determined using the method described by [Lowry et al. \(1951\)](#). The assay is based on the reaction of peptide bonds with copper (II) ions under alkaline conditions (biuret reaction), followed by reduction of the Folin–Ciocalteu reagent by the copper–protein complex, producing a blue colour whose intensity is proportional to the protein concentration. Bovine serum albumin (BSA) was used as the standard protein. A series of standard solutions of BSA (0–100 µg/mL) was prepared by serial dilution in distilled water. One milliliter of EPS solution (or standard) was mixed with 5 mL of alkaline copper solution containing 2% Na<sub>2</sub>CO<sub>3</sub> in 0.1 N NaOH, 0.5% CuSO<sub>4</sub>, and 1% sodium potassium tartrate. The mixture was incubated at room temperature for 10 min, after which 0.5 mL of freshly diluted Folin–Ciocalteu reagent (1N) was rapidly added with mixing. The reaction mixtures were allowed to stand at room temperature for 30 min for colour development.

The absorbance of the developed colour was measured at 750 nm using a UV–Visible spectrophotometer against a reagent blank. A standard calibration curve was constructed by plotting absorbance against BSA concentration, and the protein content of EPS samples was determined from the curve and expressed as µg/mL.

#### *Determination of carbohydrate content of EPS*

The carbohydrate content of the exopolysaccharide (EPS) was estimated using the phenol–sulfuric acid method ([Dubois et al., 1956](#)). The principle of this method is based on the dehydration of carbohydrate molecules by concentrated sulfuric acid, forming furfural derivatives that react with phenol to produce a coloured complex measurable at 490 nm. EPS was dissolved in distilled water to prepare the test solution. Glucose was used as the reference standard. One milliliter (1 mL) of EPS solution or standard glucose solution (0–100 µg/mL) was

## RESEARCH ARTICLE

pipetted into a test tube. To this, 1 mL of 5% phenol solution was added, followed immediately by 5 mL of concentrated sulfuric acid. The tubes were mixed thoroughly and allowed to stand at room temperature for 30 min for colour development. The absorbance of the reaction mixtures was read at 490 nm against a reagent blank using a UV–Visible spectrophotometer. A calibration curve was constructed by plotting absorbance values against standard glucose concentrations. The carbohydrate content of the EPS was then calculated from the curve and expressed as  $\mu\text{g}/\text{mL}$ .

***Biomass characterization******Determination of protein content of biomass***

The protein content of the mycelial biomass was determined following the procedure (Lowry *et al.*, 1951). The mycelial biomass was first homogenized and extracted in distilled water. The extract was centrifuged at  $5,000 \times g$  for 10 min to remove insoluble materials, and the supernatant was collected as the protein extract. Protein content in the mycelial biomass extract was expressed as  $\mu\text{g}/\text{mL}$ .

***Determination of Carbohydrate Content of Biomass***

The carbohydrate content of the mycelial biomass was determined using the phenol–sulfuric acid method (Dubois *et al.*, 1956). Carbohydrate content in the biomass extract was expressed as  $\mu\text{g}/\text{mL}$ .

***Statistical analysis***

All experiments were conducted in triplicate, and results are expressed as mean  $\pm$  standard error (SE). Statistical analysis was performed using two-way analysis of variance (ANOVA). Where significant effects were detected, Tukey's post hoc test was applied to identify differences among means. Statistical analyses were carried out using Microsoft Excel (version 2019, Microsoft Corporation, Redmond, WA, USA), and differences were considered statistically significant at  $p < 0.05$ .


**3. Results and discussion*****Growth and exopolysaccharide production of *L. squarrosulus****

The temporal patterns of exopolysaccharide (EPS) production and fungal biomass accumulation of *L. squarrosulus* cultured in Potato dextrose broth (PDB) and Yeast extract broth (YEB) over a 21-day fermentation period are presented in Figure 1a and 1b, respectively. In PDB, EPS


## RESEARCH ARTICLE

concentration increased progressively up to day 9, reaching a maximum of  $0.46 \pm 0.01$  mg/mL, followed by a gradual decline toward day 21. A similar trend was observed for biomass accumulation, which peaked at  $6.83 \pm 0.04$  mg/mL on day 9 before decreasing thereafter. In contrast, YEB supported comparatively lower and more variable EPS and biomass production, characterized by an early increase up to day 9, a decline by day 15, and a delayed secondary peak on day 18 for both parameters.

Statistical analysis revealed that culture medium, fermentation time, and their interaction significantly influenced EPS production and biomass accumulation ( $p < 0.05$ ), confirming distinct metabolic responses of *L. squarrosulus* in the two media. The higher EPS yield and biomass accumulation observed in PDB are likely attributable to its high carbohydrate content, which provides readily available carbon sources that support sustained mycelial growth and facilitate the redirection of excess carbon toward polysaccharide synthesis after the exponential growth phase (Kadam *et al.*, 2022). Conversely, YEB is predominantly nitrogen-rich with comparatively lower fermentable carbohydrate content, resulting in fluctuating biomass formation and intermittent EPS accumulation. The delayed secondary peaks observed in YEB may reflect metabolic adaptation or utilization of alternative nutrient reserves during later stages of fermentation, as previously reported in submerged fungal cultures under nutrient-limited conditions (Yau *et al.*, 2024).



a)



b)

**Fig. 1.** Growth and exopolysaccharide production of *L. squarrosulus* during submerged fermentation. (a) Exopolysaccharide (EPS) concentration (mg/mL) and (b) fungal biomass (mg/mL) measured over a 21-day fermentation period in Potato Dextrose Broth (PDB) and Yeast Extract Broth (YEB). Values represent mean  $\pm$  standard error ( $n = 3$ ). Two-way ANOVA showed significant effects of culture medium, fermentation time, and their interaction on EPS production and biomass accumulation ( $p < 0.05$ ). PDB, Potato Dextrose Broth; YEB, Yeast Extract Broth.


#### **pH dynamics in PDB and YEB media during 21-day fermentation**

To evaluate the metabolic activity of *L. squarrosulus* during submerged fermentation, pH changes in PDB and YEB were monitored over a 21-day period (Figure 2). In PDB, the initial pH of  $5.60 \pm 0.02$  at day 0 declined steadily to a minimum value of  $2.83 \pm 0.02$  on day 18, followed by a slight increase to  $2.90 \pm 0.08$  on day 21. In contrast, YEB exhibited greater pH variability, beginning with a higher initial pH of  $6.31 \pm 0.02$ , decreasing sharply to  $4.50 \pm 0.04$  by day 3, increasing slightly to  $5.09 \pm 0.02$  on day 6, and subsequently declining to its lowest pH of  $3.00 \pm 0.07$  on day 18 before rising marginally to  $3.14 \pm 0.04$  on day 21. Statistical analysis revealed that culture medium, fermentation time, and their interaction significantly influenced pH changes during fermentation ( $p < 0.05$ ).

The progressive acidification observed in PDB suggests active carbohydrate metabolism by *L. squarrosulus*, where the utilization of dextrose and potato-derived sugars likely resulted in the accumulation of organic acids. This metabolic behavior is commonly associated with enhanced fungal activity and has been reported to accompany exopolysaccharide (EPS) production during submerged fermentation (Sepúlveda *et al.*, 2025). Dynamic pH changes are known to influence

## RESEARCH ARTICLE

fungal growth performance and metabolite synthesis, including antioxidant activity, which aligns with the trends observed in this study. In contrast, the fluctuating pH profile observed in YEB may be attributed to its nutrient composition, which is rich in amino acids, peptides, and vitamins but relatively low in fermentable carbohydrates. Additionally, the intrinsic buffering capacity of yeast extract may limit drastic pH shifts, thereby reducing acidogenic metabolism and contributing to the observed variability in pH changes (Suyoon *et al.*, 2024).



**Fig. 2.** pH profile of *L. squarrosulus* culture media during fermentation. This line graph illustrates the changes in pH levels measured over a 21-day period for *L. squarrosulus* cultures grown in Potato Dextrose Broth (PDB) and Yeast Extract Broth (YEB). Values represent mean  $\pm$  standard error ( $n = 3$ ). Statistical analysis was performed using two-way ANOVA, with significance set at  $p < 0.05$ . PDB, Potato Dextrose Broth; YEB, Yeast Extract Broth.

#### FT-IR profile of exopolysaccharide


The structural features of the extracted exopolysaccharide were examined using FT-IR spectroscopy, and the resulting FT-IR spectrum is presented in Figure 3. The spectrum indicated absorption bands at the following wavenumbers and corresponding absorbance strength: 3277.4  $\text{cm}^{-1}$  (broad, intense), 2921.8  $\text{cm}^{-1}$  (medium), 2852.4  $\text{cm}^{-1}$  (medium), 1637.4  $\text{cm}^{-1}$  (moderate), 1544.0  $\text{cm}^{-1}$  (weak–moderate), 1368.5  $\text{cm}^{-1}$  (medium), 1313.7  $\text{cm}^{-1}$  (weak), 1204.0  $\text{cm}^{-1}$  (medium), 1027.3  $\text{cm}^{-1}$  (strong), and 893.4  $\text{cm}^{-1}$  (sharp, weak). The FT-IR spectrum revealed multiple characteristic absorption bands across the mid-infrared region, providing the spectral fingerprint of the extracted exopolysaccharide. Peaks at 2921.8  $\text{cm}^{-1}$  and 2852.4  $\text{cm}^{-1}$  correspond to carbon-hydrogen (C-H) stretching vibrations of aliphatic methyl and methylene groups, which

## RESEARCH ARTICLE

are indicative of sugar chains in exopolysaccharides (Patel *et al.*, 2023). A broad peak observed at  $3277.37\text{ cm}^{-1}$  is indicative of hydroxyl (O-H) stretching vibrations, which shows a high density of O-H bonds common to carbohydrate molecules and polyhydroxilic compounds contributing to their hydrophilic nature (Bhandary and Alagesan, 2024).

A moderate peak at  $1637.40\text{ cm}^{-1}$  can be associated with carbonyl (C=O) stretching (Bhandary and Alagesan, 2024). This peak may also indicate bending vibrations of uronic acid groups commonly associated with hydrated microbial EPS (Chen *et al.*, 2023).

The absorption band at  $1368.5\text{ cm}^{-1}$  corresponds to C-H bending or symmetric stretching of  $\text{COO}^-$  groups, which is often present in acidic polysaccharides with a high content of uronic acid (Yang *et al.*, 2024). The peaks at  $1313.7\text{ cm}^{-1}$  and  $1544\text{ cm}^{-1}$  may indicate amide II and amide I regions, indicating trace levels of proteinaceous components which might have been co-extracted with EPS (Sandt, 2024). The band at  $893.4\text{ cm}^{-1}$  indicates the presence of  $\beta$ -glycosidic linkages between the monosaccharide units, a common configuration in exopolysaccharides (Chen *et al.*, 2023). A notable strong band at  $1027.25\text{ cm}^{-1}$  was observed, which corresponds to C–O stretching vibrations of glycosidic linkages and pyran ring configuration, which is a key structure and a characteristic peak of polysaccharide molecules (Huang *et al.*, 2025).



**Fig. 3.** FTIR Spectrum of exopolysaccharides (EPS) produced from *L. squarrosulus*. This spectrum illustrates the percentage transmittance (%T) across a range of wavenumbers ( $\text{cm}^{-1}$ ) showing characteristic

## RESEARCH ARTICLE

absorption bands corresponding to functional groups present in the EPS. Key peaks include  $3277.37\text{ cm}^{-1}$  (O–H stretching),  $2921.8$  and  $2852.4\text{ cm}^{-1}$  (C–H stretching),  $1637.4\text{ cm}^{-1}$  (amide I, C=O stretching),  $1544\text{ cm}^{-1}$  (amide II, N–H bending),  $1368.5\text{ cm}^{-1}$  (C–H bending),  $1313.7\text{ cm}^{-1}$  (C–O stretching),  $1204$  and  $1027.25\text{ cm}^{-1}$  (C–O–C and C–O stretching of polysaccharides), and  $893.4\text{ cm}^{-1}$  (anomeric region indicating  $\beta$ -glycosidic linkages).

### **Water solubility and water absorption capacity of exopolysaccharide**

The water solubility index and water absorption capacity of the crude EPS are shown in Table 1. The crude EPS exhibited a WSI of  $54.55 \pm 0.80\%$  and a WAC of  $91.40 \pm 1.20\%$ . The extracted EPS showed moderate solubility in water and a high ability to absorb water. The water solubility of the exopolysaccharide ( $>40\%$ ) is consistent with previous reports for EPS produced ([Ogunremi and Oriyomi, 2025](#)) and aligns with the properties of fungal exopolysaccharides (EPS), which often consist of mannans and water-soluble  $\beta$ -glucans ([Amanda et al., 2024](#)). While the remaining insoluble fraction may be made up of complex or high-molecular-weight polysaccharides that are less likely to dissolve, the solubility indicates that a sizable amount of the EPS dissolves in water, most likely as a result of hydrophilic polysaccharide structures ([Ben Akacha et al., 2023](#)). The high-water absorption capacity (WAC) shows that the EPS can absorb water at a rate of  $91.40\%$  of its weight. Fungal polysaccharides, especially  $\beta$ -glucans, have a great ability to store water and create gel-like structures by hydrogen bonding ([Ogunremi and Oriyomi, 2025](#)). With respect to the WAC, the EPS possesses significant hydration qualities, which increase its potential for use in medicines, food, and cosmetics.

**Table 1.** Functional properties of exopolysaccharides from *L. squarrosulus*

Parameter	Value (%)
Water Solubility Index (WSI)	$54.55 \pm 0.80$
Water Absorption Capacity (WAC)	$91.40 \pm 1.20$

*Values are mean  $\pm$  standard error (n = 3)*

### **Antioxidant activity of exopolysaccharide**

The antioxidant capacity of the extracted EPS was determined using the DPPH radical scavenging assay, and the values are presented in Table 2. The EPS sample at  $0.016\text{ mg/mL}$  gave an absorbance of  $0.034$ , which corresponds to a percentage DPPH radical scavenging activity of  $87.41 \pm 1.05\%$ . The antioxidant activity of the EPS sample was equivalent to  $0.415 \pm 0.004\text{ mg/mL}$  of gallic acid using the equation generated from the trendline of the gallic acid standard

## RESEARCH ARTICLE

curve ( $y = 14.897x + 81.222$ ). The EPS showed a high DPPH radical scavenging activity at low concentration, with equivalent antioxidant strength comparable to gallic acid.

The antioxidant activity of the EPS from this study demonstrated a strong free radical scavenging potential. At a concentration of 0.016 mg/mL, the EPS achieved an inhibition of 87.41%, which corresponds to a gallic acid equivalent of approximately 0.415 mg/mL. This level of inhibition is notably attained at a lower concentration than that required for gallic acid to produce a comparable effect. The observed antioxidant activity is possibly due to phenolic structures and hydroxyl groups that are able to donate hydrogen atoms or electrons (Chotmanee *et al.*, 2025). Furthermore, the functional groups and monosaccharide compositions of EPS are prominently suggested to contribute to their antioxidant activity (Bouzaiene *et al.*, 2024).

**Table 2.** Antioxidant activity of exopolysaccharides from *L. squarrosulus*

Sample	Concentration (mg/mL)	DPPH scavenging activity (%)	GAE (mg/mL)
EPS	0.016	87.41 ± 1.05	0.415 ± 0.004

Values are mean ± standard error (n = 3). GAE = gallic acid equivalent EPS = exopolysaccharides

### **Protein and carbohydrate content of mycelial biomass and exopolysaccharides**

The protein and carbohydrate compositions of the mycelial biomass and EPS are shown in Table 3. While the exopolysaccharide showed  $24.34 \pm 0.44\%$  and  $62.26 \pm 0.16\%$  protein and carbohydrate content, respectively. YEB biomass gave a protein content of  $42.91 \pm 0.41\%$  and carbohydrate content of  $49.04 \pm 0.84\%$ . and PDB biomass showed a protein content of  $17.03 \pm 0.04\%$  and carbohydrate content of  $32.84 \pm 1.02\%$ . The EPS exhibited the highest carbohydrate proportion, whereas biomass produced in YEB contained the highest protein content.

The compositional analysis of cellular biomass and exopolysaccharides (EPS) revealed notable differences in protein and carbohydrate distribution depending on the growth medium. The EPS fraction was predominantly carbohydrate-rich, with 62.26% carbohydrate and 24.34% protein, consistent with the biochemical nature of EPS as high-molecular-weight sugar polymers often conjugated with small amounts of proteins or glycoproteins. This carbohydrate-rich profile of EPS aligns with other reports, which described EPS as predominantly polysaccharide in nature (Mouro *et al.*, 2024). In contrast, the cellular biomass composition varied substantially between media. Biomass derived from YEB medium exhibited higher protein content (42.91%) compared

## RESEARCH ARTICLE

to carbohydrate (49.04%), reflecting the nitrogen-rich nature of YEB that promotes protein synthesis and structural macromolecule formation. Conversely, Potato Dextrose Broth (PDB) biomass contained lower protein levels (17.03%) and a more moderate carbohydrate fraction (32.84%), suggesting a metabolism directed primarily towards carbohydrate utilization and energy storage rather than protein accumulation.

**Table 3.** Protein and carbohydrate content of mycelial biomass and exopolysaccharide from *L. squarrosulus*

Sample	Protein Content (%)	Carbohydrate Content (%)
EPS	24.34 ± 0.44	62.26 ± 0.16
Biomass in YEB	42.91 ± 0.41	49.04 ± 0.84
Biomass in PDB	17.03 ± 0.04	32.84 ± 1.02

Values are mean ± standard error (n = 3). EPS = Exopolysaccharides, YEB= Yeast Extract Broth. PDB = Potato Dextrose Broth

#### 4. Conclusions

This study demonstrated the successful production and characterisation of exopolysaccharides (EPS) and biomass from *L. squarrosulus* using Potatoes Dextrose Broth (PDB) and Yeast Extract Broth (YEB) under submerged fermentation. PDB supported higher EPS and biomass compared to YEB, with notable variations over the 21 days. FT-IR analysis confirmed the polysaccharide nature of the EPS. The crude EPS showed good water absorption capacity and strong antioxidant activity. These findings suggest *L. squarrosulus* EPS possesses desirable structural and functional properties with potential pharmaceutical applications.

**Acknowledgments.** The authors acknowledge the technical assistance provided by the laboratory technologists and the administrative support given by the department. We also acknowledge the assistance from colleagues who contributed to routine analysis.

#### References

1. Abd-Alla M.H., Hassan E.A., Mohammed E.A., Khalaf D.M., Bashandy S.R. (2025) Bacterial exopolysaccharides: a comprehensive review on sustainable production from spoiled fruit wastes. *Biocatalysis and Agricultural Biotechnology*, 66, 103607. <https://doi.org/10.1016/j.biab.2025.103607>

## RESEARCH ARTICLE

2. Asonibare A.O., Olasupo O.O., Osunlaja O.A., Oladipo A.D., Olaoti-Laaro S.O., Abdazeez A.O. (2023) Effects of mineral composition of different wood wastes in Ibadan on proximate composition of *L. squarrosulus* (Mont.). *Journal of Applied Sciences and Environmental Management*, 27, 1413–1416. <https://doi.org/10.4314/jasem.v27i7.12>
3. Ben Akacha B., Michalak M., Najar B., Venturi F., Taglieri I., Kačániová M. *et al.* (2023) Recent advances in the incorporation of polysaccharides with antioxidant and antibacterial functions to preserve the quality and shelf life of meat products. *Foods*, 12, 1647. <https://doi.org/10.3390/foods12081647>
4. Bhandary T., Alagesan P.K. (2024) Assessment of bioactivity of the novel exopolysaccharide secreted by *Bacillus subtilis* isolated from the gut of marine anchovies. *Journal of Applied Biology and Biotechnology*, 12, 205–212.
5. Bouzaiene T., Vall M., Ziadi M., Rejeb I., Yangui I., Abdelkarim A. *et al.* (2024) Exopolysaccharides from *Lactiplantibacillus plantarum* C7 exhibited antibacterial, antioxidant, anti-enzymatic, and prebiotic activities. *Fermentation*, 10, 339. <https://doi.org/10.3390/fermentation10070339>
6. Chen C.C., Nargotra P., Kuo C.H., Liu Y.C. (2023) High-molecular-weight exopolysaccharides production from *Tuber borchii* cultivated by submerged fermentation. *International Journal of Molecular Sciences*, 24, 4875. <https://doi.org/10.3390/ijms24054875>
7. Chotmanee T., Suwannarach N., Kumla J., Phongphisutthinant R., Chaipoot S., Wiriyacharee P., Tantibhadrasapa A., Li S., Thiennimitr P., Lumyong S. (2025) Exopolysaccharide production by seven basidiomycetous fungi and their antioxidant and immunomodulatory activities against *Salmonella* infection. *Frontiers in Cellular and Infection Microbiology*, 15, 1610403. DOI: 10.3389/fcimb.2025.1610403.
8. Dubois M., Gilles K.A., Hamilton J.K., Rebers P.A., Smith F. (1956) Colorimetric method for determination of sugars and related substances. *Analytical Chemistry*, 28, 350–356.
9. Fischer A., Castagna V., Omer S., Marmorstein M., Wu J., Ceballos S., Skoog E., Lebrilla C.B., Suarez C., Schnitzler A. (2024) Characterization of the exopolysaccharides produced by the industrial yeast *Komagataella phaffii*. *Journal of Industrial Microbiology and Biotechnology*, 51, kuae046. DOI: 10.1093/jimb/kuae046.
10. Gyamfi M.A., Yonamine M., Aniya Y. (1999) Free radical scavenging action of medicinal herbs from Ghana: *Thonningia sanguinea* on experimentally induced liver injuries. *General Pharmacology*, 32, 661–667.

## RESEARCH ARTICLE

11. Hibbett D.S., Binder M., Bischoff J.F., Blackwell M., Cannon P.F. *et al.* (2007) A higher-level phylogenetic classification of the Fungi. *Mycological Research*, 111, 509–547. <https://doi.org/10.1016/j.mycres.2007.03.004>
12. Huang Y., Liang W., Lu Y., Xiong J., Liu D., Jia X. (2025) Identification, antioxidant and immunomodulatory activities of a neutral exopolysaccharide from *Lactiplantibacillus plantarum* DMDL 9010., *Nutrients*, 17(14), 2265. <https://doi.org/10.3390/nu17142265>
13. Kadam S., Karne S., Maybhate M., Maniyar A., Gore A., Mehta H. (2022) Support by potato dextrose agar for fungal and bacterial growth at 40°C. *International Journal of Microbial Science*, 3.
14. Kiran N.S., Yashaswini C., Singh S., Prajapati B.G. (2024) Revisiting microbial exopolysaccharides: a biocompatible and sustainable polymeric material for multifaceted biomedical applications. *3 Biotech*, 14, 95.
15. Kumla J., Suwannarach N., Sujarit K., Penkhruue W., Kakumyan P., Jatuwong K., Vadhanarat S., Lumyong S. (2020) Cultivation of mushrooms and their lignocellulolytic enzyme production through the utilization of agro-industrial waste. *Molecules*, 25, 2811. <https://doi.org/10.3390/molecules25122811>
16. Lee S., Kim E., Jo M., Choi Y.J. (2024) Characterization of yeast protein isolates extracted via high-pressure homogenization and pH shift: A promising protein source enriched with essential amino acids and branched-chain amino acids., *Journal of Food Science*, 89, e16918. <https://doi.org/10.1111/1750-3841.16918>
17. Lu Y., Qin Y., Sun Y., Joseph C.M.L., Bisson L.F., Song Y., Liu Y. (2024) Insight into the growth and metabolic characteristics of indigenous commercial *Saccharomyces cerevisiae* NX11424 at high and low levels of yeast assimilable nitrogen based on metabolomic approach., *Food Microbiology*, 124, 104593. <https://doi.org/10.1016/j.fm.2024.104593>
18. Lowry O.H., Rosebrough N.J., Farr A.L., Randall R.J. (1951) Protein measurement with the Folin phenol reagent. *Journal of Biological Chemistry*, 193, 265–275.
19. Łysakowska P., Sobota A., Wirkijowska A. (2023) Medicinal mushrooms: their bioactive components, nutritional value and application in functional food production. *Molecules*, 28, 5393. <https://doi.org/10.3390/molecules28175393>
20. Mouro C., Gomes A.P., Gouveia I.C. (2024) Microbial exopolysaccharides: Structure, diversity, applications, and future frontiers in sustainable functional materials., *Polysaccharides*, 5(3), 241–287. <https://doi.org/10.3390/polysaccharides5030018>

## RESEARCH ARTICLE

21. Ogidi C.O., Ubaru A.M., Ladi-Lawal T., Thonda O.A., Aladejana O.M., Malomo O. (2020) Bioactivity assessment of exopolysaccharides produced by *Pleurotus pulmonarius* in submerged culture with different agro-waste residues. *Helijon*, 6, e05685.
22. Ogunremi O.R., Oriyomi O.V. (2025) Technological and functional properties of exopolysaccharides produced by *Weissella confusa* strains isolated from kunu-zaki. *Food Science and Technology International*.
23. Patel V.H., Shah K., Patel F.R. (2023) Physical characterization and structure elucidation of exopolysaccharide from marine halotolerant bacterium *Enterobacter cloacae* VHP-34. *Industrial Biotechnology*, 19. <https://doi.org/10.1089/ind.2023.0020>
24. Reddy C.K., Suriya M., Haripriya S. (2013) Physico-chemical and functional properties of resistant starch from red kidney beans (*Phaseolus vulgaris* L.) prepared by enzymatic method. *Carbohydrate Polymers*. <https://doi.org/10.1016/j.carbpol.2013.02.060>
25. Ruas-Madiedo P., Salazar N., de los Reyes-Gavilán C.G. (2010) Exopolysaccharides produced by lactic acid bacteria in food and probiotic applications. In: Holst O., Brennan P.J., von Itzstein M., Moran A.P. (eds.) *Microbial Glycobiology*, Academic Press, 885–902.
26. Salimi F., Farrokh P. (2023) Recent advances in the biological activities of microbial exopolysaccharides. *World Journal of Microbiology and Biotechnology*, 39, 213.
27. Sandt C. (2024) Identification and classification of proteins by FTIR microspectroscopy: A proof of concept., *Biochimica et Biophysica Acta – General Subjects*, 1868(10), 130688. DOI: 10.1016/j.bbagen.2024.130688.
28. Sepúlveda Del Rio Hamacek H., Tingajeva O., Ostertag K., Reķēna A., Illarionov A., Jõul P., Monteiro de Oliveira P., Martín-Hernández G.L.C., Müller B., Bonturi N., Passoth V., Lahtvee P., Kumar R. (2025) Acidification by nitrogen metabolism triggers extracellular biopolymer production in an oleaginous yeast., *Applied and Environmental Microbiology*, 91, e00947-25. DOI: 10.1128/aem.00947-25.
29. Stabnikova O., Stabnikov V., Paredes-Lopez O. (2024) Wild and cultivated mushrooms as food, pharmaceutical and industrial products. *Ukrainian Food Journal*, 13, 20–59.
30. Synytsya A., Novák M. (2014) Structural analysis of glucans. *Annals of Translational Medicine*, 2, 17.

**RESEARCH ARTICLE**

31. Wang C.S., Virgilio N., Wood-Adams P.M., Heuzey M.C. (2018) A gelation mechanism for gelatin/polysaccharide aqueous mixtures. *Food Hydrocolloids*, 79, 462–472. <https://doi.org/10.1016/j.foodhyd.2018.01.016>

32. Wang W., Ju Y., Liu N., Hao L. (2023) Structural characteristics of microbial exopolysaccharides in association with their biological activities. *Chemical and Biological Technologies in Agriculture*, 10.

33. Yang R., Liu L., Gao D., Zhao D. (2024) Purification, structural characterization, and bioactive properties of exopolysaccharides from *Saccharomyces cerevisiae* HD-01. *Frontiers in Bioengineering and Biotechnology*, 12, 1455708. <https://doi.org/10.3389/fbioe.2024.1455708>

## RESEARCH ARTICLE

## SUPPLEMENTARY MATERIAL

Supplementary Table S1. FT-IR fingerprints of EPS from *L. squarrosulus*

Wavenumber (cm <sup>-1</sup> )	Transmittance (%T)	Functional Group / Vibration	Functional Property	Reference
893.4	~90%	β-Glycosidic linkage (C–H deformation)	Indicates β-configuration of sugar monomers	(Mouro <i>et al.</i> , 2024)
1027.25	~88%	C–O stretching (pyranose ring)	Core polysaccharide structure / fingerprint region	(Mouro <i>et al.</i> , 2024)
1204.0	~94%	C–O–H or C–O–C vibration	Glycosidic linkage in sugar rings	(Mouro <i>et al.</i> , 2024)
1313.7	~94%	C–N stretch / CH <sub>2</sub> wag	Amine group, potential nitrogen-containing residues	(Chen <i>et al.</i> , 2023)
1368.5	~93%	C–H bending / COO <sup>-</sup> symmetric stretch	Uronic acid presence / acidic polysaccharide marker	(Yang <i>et al.</i> , 2024)
1544.0	~93%	Amide II (N–H bending)	Possible protein residue or amine contamination	(Chen <i>et al.</i> , 2023)
1637.40	~92%	C=O stretching / O–H bending	Carboxylic groups, water content	(Chen <i>et al.</i> , 2023)

## RESEARCH ARTICLE

2852.4	~91%	C–H	symmetric stretching (–CH <sub>3</sub> )	Structural aliphatic groups	(Mouro <i>et al.</i> , 2024)
2921.8	~91%	C–H	asymmetric stretching (–CH <sub>2</sub> )	Backbone sugar rings	(Mouro <i>et al.</i> , 2024)
3277.37	~89% (estimate)	O–H (broad)	stretching	Hydrogen bonding, hydrophilicity	(Mouro <i>et al.</i> , 2024)